Spinal cord - repetition





© David Kachlík 30.9.2015

Ascending tracts

- tractus spino-bulbo-thalamo-corticalis

= posterior fascicle tract = lemniscal system = fasciculus gracilis + cuneatus

  – *propriception, fine skin sensitivity, discrimination feeling, push, pressure, vibration*

- tractus spinothalamicus ant.+lat. = anterolateral system
  – *Fast pain, warm + cold, rough skin sensitivity*
- tractus spinoreticularis
  – *Slow pain*
- tractus spinocerebellares
- ant.+post.
- *And other……☺*



© David Kachlík 30.9.2015

Descending tracts

- tractus corticospinalis = pyramidal tract
  - Principal motor tract
  - 1st neuron – cortex (Betz pyramidal cell)
  - 2nd neuron – alfa-moto neuron → spinal nerve
- Extrapyramidal trycts
  - tr. reticulospinalis – gamma moto-neurons
  - tr. vestibulospinalis – postural muscles
  - tr. rubrospinalis (rudimentary)
  - other ☺



© David Kachlík 30.9.2015

BRAIN STEM

David Kachlík
Petr Zach

© David Kachlík 30.9.2015

Brain stem - *function*

- Convertor of all převaděč všech ***ascending and descending tracts***

- **Reticular formation (RF)**

Life important ***reflex centers*** – heart activity, breathing, vazomotorics, consciousness

- Nuclei of cranial nerves n. III - XII

© David Kachlík 30.9.2015

Brain stem (Truncus encephali)

truncus encephali

- **medulla oblongata** = oblongate
- **pons** (Varoli) = pons
- **mesencephalon** = mid brain

Cavities of brain stem

- **ventriculus quartus** = 4th ventricle
  - fossa rhomboidea = base of 4th ventricle
- **aqueductus mesencephali** (Sylvii) = Sylvius canal, aqueduct

© David Kachlík 30.9.2015



© David Kachlík 30.9.2015





© David Kachlík 30.9.2015

Medulla oblongata = My
medullae spinal

- Cranial continuation of s

Ventral side:

- ***pyramis***
  - Fibers of tractus corticosp
  - decussatio pyramidum (cr
- ***oliva***
  - complexus olivaris inferior
  - Ventrally exit n.XII (sulcus
  - Dorsally exit n.IX, X, XI (s
- ***tuberculum trigeminal***
  - ncl. + tractus spinalis n.V.



© David Kachlík 30.9.2015

Pons and oblongate ventral view



1 - n. IX.
2 - n. X.
3 - radix spinalis n. XI.
4 - radix cranialis n. XI.
5 - sulcus basilaris
6 - n. VI.
7 - pars motoria n. V.
8 - pars sensoria n. V.
9 - trigeminofacial line (Henlei)
10 - pedunculus cerebellaris medius
11 - n. VII.
12 - n. intermedius
13 - n. VIII.
14 - pedunculus cerebellaris inferior
15 - foramen caecum medullae oblongatae
16 - n. XII
17 - decussatio pyramidum
18 - radix anterior C1

© David Kachlík 30.9.2015





© David Kachlík 30.9.2015

Medulla oblongata = Myelencephalon = Bulbus medullae spinalis = Oblongate

Dorsal side:

- ***tuberculum gracile + cuneatum***

tractus bulbo-spinalis (fasciculus gracilis Golli + cuneatus Burdachi) → **ncl. gracilis + cuneatus** → fibrae arcuatae internae → decussatio lemnisci medialis → lemniscus medialis

- ***pedunculi cerebellares inferiores***

continue as funiculus lateralis of spine + ncl. cuneatus accessorius → fibrae arcuatae externeae posteriores

→ corpus restiforme

→ corpus juxtarestiforme

→ carry tracts to cerebellum

© David Kachlík 30.9.2015

Oblongate – internal structure and nuclei

- complexus olivaris inferior (3 nuclei)
- ncl. gracilis, cuneatus, cuneatus accessorius
- Nuclei of cranial nerves
  - n. V, IX, X, XI, XII
- Nuclei of RF
  - nuclei reticulares, raphes
- ncl. arcuati – connected same way as ncll. pontis

© David Kachlík 30.9.2015

Oblongate – internal structure and tracts

- tr. corticospinalis (ventrally)
- tr. spinocerebellaris ant., post., rostralis
- tr. spino-bulbaris → lemniscus medialis
- tr. spinothalamicus ant.+lat. → lemniscus spinalis
- tr. spinoreticularis → RF
- fasciculus longitudinalis medialis + posterior
- Extrapyramidal tracts
  - tr. reticulospinalis, vestibulospinalis, tectospinalis, rubrospinalis, interstitiospinalis
  - raphaespinalis, caerulospinalis

© David Kachlík 30.9.2015

Medulla - sectio in decussatione pyramidum



© David Kachlík 30.9.2015

Bottom of the fourth ventricle

(FOSSA RHOMBOIDEA)



1 - lingula cerebelli
2 - sulcus medianus
3 - pedunculus cerebellaris superior
4 - pedunculus cerebellaris medius
5 - pedunculus cerebellaris inferior
6 - locus caeruleus
7 - sulcus limitans
8 - colliculus facialis
9 - area vestibularis
10 - striae medullares ventriculi quarti
11 - taenia cinerea
12 - tuberculum trigeminale
13 - trigonum n. X.
14 - trigonum n. XII.
15 - tuberculum cuneatum
16 - tuberculum gracile
17 - eminentia medialis
18 - fasciculus gracilis /Golli/
19 - fasciculus cuneatus /Burdachi/

A - section at colliculus superior
B - section at tuberculum trigeminale
C - section at tuberculum gracile et cuneatum

© David Kachlík 30.9.2015

Section of oblongate at the level of

TUBERCULUM CUNEATUM A GRACILE



1 - tuberculum gracile
2 - tuberculum cuneatum
3 - tractus spinocerebellaris posterior
4 - tractus spinocerebellaris anterior
5 - canalis centralis
6 - nucleus n. XII
7 - nucleus gracilis
8 - nucleus cuneatus
9 - nucleus spinalis n. V
10 - nucleus ambiguus
11 - tractus tectospinalis
12 - tractus rubrospinalis
13 - tractus spinothalamicus
14 - tractus corticospinalis

© David Kachlík 30.9.2015

Medulla - sectio in deccussatione lemniscorum



Fasciculus gracilis
Nucleus gracilis
Central canal
Fasciculus cuneatus
Nucleus cuneatus
Hypoglossal nucleus
Internal arcuate fibers
Spinal tract of trigeminal nerve
Medial longitudinal fasciculus
Spinal nucleus of trigeminal nerve
Posterior spinocerebellar tract
Spinal root of accessory nerve
Inferior olivary nucleus
Lateral spinothalamic tract
Anterior spinocerebellar tract
Hypoglossal nerve
Pyramid
Decussation of medial lemnisci

© David Kachlík 30.9.2015

Section of oblongate at the level of

TUBERCULUM TRIGEMINALE



1 - ventriculus quartus
2 - velum medullare superius
3 - tractus spinocerebellaris posterior
4 - tractus spinocerebellaris anterior
5 - nucleus et tractus spinalis n. V
6 - nuclei vestibulares
7 - nuclei tractus solitarii
8 - nucleus salivatorius inferior
9 - nucleus ambiguus
10 - nucleus n. XII
11 - fasciculus longitudinalis medialis
12 - tractus tectospinalis
13 - tractus rubrospinalis
14 - tractus spinothalamicus
15 - lemniscus medialis
16 - nuclei olivares
17 - tractus corticospinalis

© David Kachlík 30.9.2015

Medulla oblongata - sectio in trigone nervi hypoglossi



© David Kachlík 30.9.2015

Medulla oblongata - sectio in complexo olivare inferiore



© David Kachlík 30.9.2015

Syndroma medullaris laterale (Wallenbergi)



© David Kachlík 30.9.2015

Syndroma medullare laterale *Wallenbergi*



Lesion of **<u>a. cerebelli posterior inferior</u>**

- dysphagia + dysarthria ipsilateralis (*ncl. ambiguus*)
- analgesia + thermoanaesthesia capitis ipsilateralis (*ncl. + tractus spinalis n. V*)
- vertigo, nausea, vomitus, nystagmus (*ncll. vestibulares*)
- syndroma Claude Bernard-Horner ispilaterale (*descending sympatetic fibers*)
- Ipsilateral lesion of cerebellum
- analgesia + thermoanaesthesia contralateralis corporis (*tractus spinothalamicus – lemniscus spinalis*)

© David Kachlík 30.9.2015

Syndroma medullare mediale (Dejerinei)



© David Kachlík 30.9.2015

Syndroma medullare mediale *Dejerine*

Lesion of **<u>a. vertebralis</u>**

- paresis contralateralis (*tractus pyramidalis*)
- contralateral lesion of fine sensitivity and proprioception (*lemniscus medialis – tr. bulbothalamocorticalis*)
- hemiglossoplegia ipsilateralis (*n. XII*)



© David Kachlík 30.9.2015

Clinical syndromes of caudal half of oblongate

A. Hemiplegia alternans inf.

B. Avelis syndrom

C. Jackson syndrom

D. Tapia syndrom

1. tractus pyramidalis

2. n. hypoglossus

3. spinothalamic tract

4. nucleus ambiguus



© David Kachlík 30.9.2015

Hemiglossoplegia



© David Kachlík 30.9.2015

PONS

© David Kachlík 30.9.2015

Pons = Po

Located only ventrally

Dorsally – bottom of 4th ve
cerebellum

sulcus bulbopontinus

- Exit of n. VI

- sulcus basilaris (non pai
- pedunculi cerebellares m
  - Tracts from ncll. pontis int
- linea trigeminofacialis *He*
  - Between exits of n.V and
    cerebellum
- angulus pontocerebellari
  - Exit of n. VII and n. VIII +



© David Kachlík 30.9.2015

Pons – internal structure and parts

- pars basilaris pontis (= pars anterior; ventralis)
  - Evolutionary younger
  - Ventral part at skull base
  - nuclei pontis + descending tracts (fibrae pontis)
- tegmentum pontis (= pars posterior; dorsalis)
  - Evolutionary older
  - Dorsal part at the base of the 4th ventricle
  - Nuclei of cranial nerves
  - RF
  - Ascending tracts (lemnisci)
  - Old tracts (fasciculus longitudinalis medialis + posterior)

© David Kachlík 30.9.2015

Pons – internal structure and nuclei

- ncll. pontis
  - Connection of tract from cortex to cerebellum (tr. corticocerebellaris)
- ncl. olivaris superior
  - Nuceli connected into auditory path
  - Recognition of sound source position
- corpus trapezoideum (nuclei c.t.)
  - Nuclei connected into auditory path
- ncll. parabrachiales
  - Part of limbic system
  - Respiration, pain, taste, CTA, *serotonin*
- Nuclei of cranial nerves
  - n. V, VI, VII, VIII

© David Kachlík 30.9.2015

Base of the 4th ventricle with cranial nerve nuclei projections



1 - colliculus superior
2 - colliculus inferior
3 - frenulum veli medullaris superioris
4 - tractus mesencephalicus n. V.
5 - pedunculus cerebellaris superior
6 - pedunculus cerebellaris medius
7 - nucleus principalis n. V.
8 - nucleus vestibularis superior /*Bechtěrev*/
9 - nucleus cochlearis posterior
10 - nucleus vestibularis medialis /*Schwalbe*/
11 - nucleus vestibularis lateralis /*Deiters*/
12 - nucleus cochlearis anterior
13 - nucleus vestibularis inferior /*Roller*/
14 + 16 - nuclei tractus solitarii
15 - tractus spinalis n.V.
17 - nucleus accessorius n. III. /*Edinger-Westphal*/
18 - nucleus n III.
19 - nucleus n. IV.
20 - nucleus motorius n. V.
21 - nucleus n. VI.
22 - nucleus salivatorius superior
23 - nucleus n. VII.
24 - nucleus salivatorius inferior
25 - nucleus posterior (dorsalis) n. X.
26 - nucleus ambiguus
27 - nucleus n. XII.
28 - nucleus n. XI. (= součást ncl. ambiguus a retroambiguus)

© David Kachlík 30.9.2015

Ncll. tractus solitarii

- Viscerosensory nuclus of side mixed system
- Every aferent information from visceral organs (+taste)
- Center for autonomic nervous system (**elementary autonomic reflexes**)
  - Deals with intake and processing of nutrients
  - Salivatory reflex, secretion of gastric or pancreatic fluids
- *More complex reactions requiring integration of larger amount of elementary autonomic reflexes or coordination of autonomic+endocrine+somatic systems*
  - *Vomitus, swallowing, coughing, breathing, orgasm, threat of freezing or overheting*
  - *Regulated from RF and more importantly from hypothalamus*
- Analogy in spinal cord = ncl. intermediomedialis
  - automatic defecation, micturia, erection

© David Kachlík 30.9.2015

Pons – internal structure and tracts

pars basilaris

- ***fibrae pontis longitudinales***
  - fribrae corticospinales = scattered pyramids → pyramis
  - fibrae corticopontinae → ncll. pontis →
- ***fibrae pontis transversae*** (= fibrae pontocerebellares) → pedunculi cerebellares medii → cerebellum

© David Kachlík 30.9.2015

Pons – internal structure and tracts

tegmentum

- lemniscus medialis
- tr. spinothalamicus → lemniscus spinalis
- tr. spinocerebellaris ant.
- tr. spinoreticularis → RF
- fasciculus longitudinalis medialis + posterior
- Extrapyramid tracts
  - tr. reticulospinalis, tectospinalis, rubrospinalis, caerulospinalis, interstitiospinalis

© David Kachlík 30.9.2015

Ventriculus quartus = 4th ventricle



- floor: fossa rhomboidea
- roof (= tegmen)
  - velum medullare sup.
  - fastigium
  - velum medullare inf.
  - apertura mediana v.q. (= foramen *Magendiei*) – non paired
  - obex (caudally)
- tela choroidea ventriculi quarti
  - plexus choroideus ventriculi quarti
- recessus lateralis
  - apertura lateralis v.q. (= foramen *Luschkae*) – paired
  - Outside looking Bochdaleki trunc from plexus choroideus
- aqueductus mesencephali *Sylvii* – into 3rd ventricle

© David Kachlík 30.9.2015

Bottom of the fourth ventricle

(FOSSA RHOMBOIDEA)



1 - lingula cerebelli
2 - sulcus medianus
3 - pedunculus cerebellaris superior
4 - pedunculus cerebellaris medius
5 - pedunculus cerebellaris inferior
6 - locus caeruleus
7 - sulcus limitans
8 - colliculus facialis
9 - area vestibularis
10 - striae medullares ventriculi quarti
11 - taenia cinerea
12 - tuberculum trigeminale
13 - trigonum n. X.
14 - trigonum n. XII.
15 - tuberculum cuneatum
16 - tuberculum gracile
17 - eminentia medialis
18 - fasciculus gracilis /Golli/
19 - fasciculus cuneatus /Burdachi/

A - section at colliculus superior
B - section at tuberculum trigeminale
C - section at tuberculum gracile et cuneatum

© David Kachlík 30.9.2015



© David Kachlík 30.9.2015

Base of the 4th ventricle with cranial nerve nuclei projections



1 - colliculus superior
2 - colliculus inferior
3 - frenulum veli medullaris superioris
4 - tractus mesencephalicus n. V.
5 - pedunculus cerebellaris superior
6 - pedunculus cerebellaris medius
7 - nucleus principalis n. V.
8 - nucleus vestibularis superior /*Bechtěrev*/
9 - nucleus cochlearis posterior
10 - nucleus vestibularis medialis /*Schwalbe*/
11 - nucleus vestibularis lateralis /*Deiters*/
12 - nucleus cochlearis anterior
13 - nucleus vestibularis inferior /*Roller*/
14 + 16 - nuclei tractus solitarii
15 - tractus spinalis n.V.
17 - nucleus accessorius n. III. /*Edinger-Westphal*/
18 - nucleus n III.
19 - nucleus n. IV.
20 - nucleus motorius n. V.
21 - nucleus n. VI.
22 - nucleus salivatorius superior
23 - nucleus n. VII.
24 - nucleus salivatorius inferior
25 - nucleus posterior (dorsalis) n. X.
26 - nucleus ambiguus
27 - nucleus n. XII.
28 - nucleus n. XI. (= součást ncl. ambiguus a retroambiguus)

© David Kachlík 30.9.2015



© David Kachlík 30.9.2015





© David Kachlík 30.9.2015

Bottom of the fourth ventricle

(FOSSA RHOMBOIDEA)



1 - lingula cerebelli
2 - sulcus medianus
3 - pedunculus cerebellaris superior
4 - pedunculus cerebellaris medius
5 - pedunculus cerebellaris inferior
6 - locus caeruleus
7 - sulcus limitans
8 - colliculus facialis
9 - area vestibularis
10 - striae medullares ventriculi quarti
11 - taenia cinerea
12 - tuberculum trigeminale
13 - trigonum n. X.
14 - trigonum n. XII.
15 - tuberculum cuneatum
16 - tuberculum gracile
17 - eminentia medialis
18 - fasciculus gracilis /Golli/
19 - fasciculus cuneatus /Burdachi/

A - section at colliculus superior
B - section at tuberculum trigeminale
C - section at tuberculum gracile et cuneatum

© David Kachlík 30.9.2015

Section of the pons at the level of

COLLICULUS FACIALIS



1 - eminentia medialis
2 - area vestibularis
3 - nucleus cochlearis posterior
4 - nucleus cochlearis anterior
5 - nucleus et tractus n. V.
6 - nucleus n. VI.
7 - nucleus n. VII.
7a - nucleus salivatorius superior
8 - nuclei vestibulares
9 - fasciculus longitudinalis medialis
10 - tractus tectospinalis
11 - lemniscus medialis
12 - tractus rubrospinalis
13 - corpus trapezoideum
14 - fibrae pontis transversae
(černé příčné čáry)
nuclei pontis (velké hnědé tečky)
fibrae pontis longitudinales
/roztříštěné svazky pyramid/
(malé červené tečky)

© David Kachlík 30.9.2015

Pons - sectio in collicule faciale



© David Kachlík 30.9.2015

Pons - sectio in nucleis trigeminalibus



© David Kachlík 30.9.2015

Locked-in syndrom

- Awake patient
- Whole kvadrupalsy + palsy of cranial nerves up from n.V
- Lesion of pars basilaris pontis = closure of **a. basilaris**
- Sometimes preserved proprioception and sensitivity from body

- "*the closest thing to being buried alive".*
- *"maladie de l'emmuré vivant", "Eingeschlossensein„, Cerebromedullospinal Disconnection, De-Efferented State, Pseudocoma, ventral pontine syndrome*

© David Kachlík 30.9.2015

Locked – in syndrom



© David Kachlík 30.9.2015

Persistent vegetative state
Locked-out syndrom

- Patient non reacting to sensory impulses
- Does not mind surrounding neiter self
- Preserved cycle of sleep and wake
- Respiration without help of respiratory devices
- Vast cortical injury or fibers connecting cortex with thalamus
- Bilateral thalamic lesion
- *„coma vigile, apalic syndrom, chronic vegetative state, permanent vegetative tate (PVS)“*

© David Kachlík 30.9.2015

Persistent vegetative state „locked-out“



© David Kachlík 30.9.2015





Peripheral „Bell“ palsy

n. VII





Central palsy

© David Kachlík 30.9.2015

Strabismus convergens – n. VI



© David Kachlík 30.9.2015

Clinical syndrome of pons

A. Millard-Gubler Syndrom
B. Raymond Syndrom
C. Foville Syndrom
D. Raymond-Cestan Syndrom
E. Marie-Poix Syndrom

1. *tractus pyramidalis*
2. *fasciculus longitudinalis med.*
3. *lemniscus medialis*
4. *tractus spinothalamicus*
5. nucleus motorius n.V
6. nucleus principalis n.V
7. *tractus spinalis n. V*
8. *pedunculus cerebellaris medius*



© David Kachlík 30.9.2015

MID BRAIN

© David Kachlík 30.9.2015



© David Kachlík 30.9.2015



© David Kachlík 30.9.2015

Mesencephalo

- Cranial part of brain ster

Ventral side:

- **fossa interpeduncular**
  - Exit of n. III
  - substantia perforata poste
    - Non paired holed plate for aa.
  - cisterna interpeduncularis
- **crura cerebri**
  - Descending motor tracts (
    - fibrae corcticospinales
    - fibrae corticonucleares
    - fibrae corticopontinae (front parietopontinae)
    - fibrae corticoreticulares



© David Kachlík 30.9.2015





© David Kachlík 30.9.2015

Mesencephalo

- Cranial part of brain ste

Dorsal side:

lamina quadrigemina =

- **colliculi superiores**
  - Nuclei from optical tract
- **colliculi inferiores**
  - Nuclei from auditory path
- On sides of **brachium c inferioris**
- **Exit of n. IV**
  - nerve crosses inside mid nervorum trochlearium)
- *cisterna quadrigeminalis*



© David Kachlík 30.9.2015





© David Kachlík 30.9.2015





© David Kachlík 30.9.2015

DORSAL VIEW OF MESENCEPHALON



1 - third ventricle
2 - stria medullaris of thalamus
3 - pulvinar
4 - habenular trigone
5 - pineal gland
6 - superior colliculus
7 - brachium of inferior colliculus
8 - lateral geniculate body
9 - inferior colliculus
10 - brachium of superior colliculus
11 - medial geniculate body
12 - frenulum of superior medullary vellum
13 - trochlear nerve
14 - lingula
15 - superior cerebellar peduncle

VENTRAL VIEW OF MESENCEPHALON



1 - optic nerve
2 - optic chiasm
3 - optic tract
4 - hypophysis
5 - infundibulum hypophysis
6 - mamillary body
7 - interpeduncular fossa
8 - oculomotor nerve
9 - cerebral crus
10 - basilar sulcus
11 - pons /*Varol´s*/

© David Kachlík 30.9.2015

Mid brain – internal composition

3 parts on section: tectum, tegmentum, crus

- **tectum mesencephali (tectal plate)**

  = dorsal thin plate with two pairs of tubercles

aqueductus mesencephali *Sylvii* – arbitrary border

- **pedunculus cerebri (brain pedicle)**

  = ventral most of mid brain

  – **tegmentum mesencephali** (= pars posterior pedunculi cerebri)
    - nuclei, some descending and all ascending tracts

  – **crus cerebri** (= pars anterior *s.* basalis pedunculi cerebri)
    - Contains only descending tracts

basis pedunculi = crus cerebri + substantia nigra

© David Kachlík 30.9.2015

Mid brain – section



© David Kachlík 30.9.2015

Mesencephalic section at the level of COLLICULLUS SUPERIOR



1 - Colliculus superior
2 - Aquaeductus mesencephali */Sylvii/*
3 - Substantia grisea centralis
4 - Nucleus n. III.
5 - Tractus tectospinalis
6 - Decussatio tegmenti posterior
7 - Lemniscus medialis
8 - Nucleus ruber
9 - Tractus rubrospinalis
10 - N. III.
11 - Substantia nigra
12 - Fibrae frontopontinae */Arnoldi/* (Tractus coticopontinus)
13 - Fibrae corticonucleares (Tractus pyramidalis)
14 - Fibrae corticospinales (Tractus pyramidalis)
15 - Fibrae occipito-, parieto-, temporopontinae (Tractus corticopontinus)

Mesencephalic section at the level of COLLICULLUS INFERIOR



1 - Aquaeductus mesencephali + substantia grisea centralis
2 - Colliculus inferior
3 - Radix mesencephalica n. trigemini
4 - Lemniscus lateralis
5 - Lemniscus medialis
6 - Tractus tectospinalis
7 - "Nucleus albus"
8 - Deccussatio pedunculorom cerebellarium superiorum
9 - Substantia nigra
10 - Fibrae frontopontinae */Arnoldi/* (Tractus coticopontinus)
11 - Fibrae corticonucleares (Tractus pyramidalis)
12 - Fibrae corticospinales (Tractus pyramidalis)
13 - Fibrae occipito-, parieto-, temporopontinae (Tractus corticopontinus)
14 - Nucleus n. IV.
15 - N. IV.

© David Kachlík 30.9.2015

Mid brain
Tectum

lamina tecti = lamina quadrigemina

colliculus superior

- Evolutionary older structures of visual system
- Its function does not correspond with vision
- <u>retinotopic</u> parcellation
- Importance for processing of fast targeted movements
- contain 7 layers (*laminae)*

© David Kachlík 30.9.2015

Mid brain
Tectum

colliculus superior

- **brachium colliculi superioris** – connection to corpus geniculatum laterale (part of metatalamus)
- AF: visual path, spine, cortex
  - Superficial layers gets side tracts from optic path
- EF:
  - Deep layers connected to crossed tractus tectospinalis – eyeball movement sync and head to visual inputs
  - tractus tectointerstitialis → ncl. *Cajali* + *Darkševiči* – acommodation and convergence
  - tractus tectonuclearis → eyeball muscles

© David Kachlík 30.9.2015

Mid brain – dorsal side



© David Kachlík 30.9.2015

Mid brain
Tectum

lamina tecti = lamina quadrigemina

colliculus inferior

- Nucleus of auditory path (3rd neuron)
  - 3 subnuclei
- laminar tonotopic organization
- **brachium colliculi inferioris**
  - Continuation of auditory path into corpus geniculatum mediale (part of metatalamus), in which is 4th neuron

© David Kachlík 30.9.2015

Diencephalon
Epithalamus
Area pretectalis

ncll. pretectales

- 4 nuclei, grants pupillary reflex
- AF: from optic tract
- EF:
  - into nucleus n. III accessorius dorsalis *Edingeri-Westphali* → *next as parasympatic path to* m. sphincter pupillae → ***miosis***
  - into RF → centrum ciliospinale *Budgei* (segments C8-T1) → truncus sympathicus → ganglion cervicale superius (*connection*) → *as sympatetic path to* m. dilatator pupillae → ***mydriasis***

© David Kachlík 30.9.2015

Mid brain
Tegmentum – nuclei
Nucleus ruber (red nucleus)

motoric nucleus at the level of colliculus superior

- ***pars magnocellularis*** (*in human and apes rudimentary)*

  → tractus rubrospinalis (crossing at decussatio tegmentalis anterior) – somatotopically in whole spine → activation of flexors

- ***pars parvocellularis***

  → fibrae rubroolivares = non crossed tracts running in the middle of tegmentum in tractus tegmentalis centralis

  – Part of *Papez* cerebellar circuit: cerebellum → NR → oliva → cerebellum

- AF: from cortex and cerebellum
- EF: into oliva and cerebellum, into RF, into talamus

© David Kachlík 30.9.2015

Mesencephalon - sectio in collicule superiore



© David Kachlík 30.9.2015

Mesencephalon - sectio in collicule inferiore



© David Kachlík 30.9.2015

Mid brain
Tegmentum – nuclei
Substantia nigra Soemmeringi

- motoric nucleus connected with mid brain
- ***pars compacta (A9)***
  - Produces dopamin
  - Makes striatum functional
  - Interposited basal ganglion
  - *In case of lack of dopamin* → ***Parkinson disease***
- ***pars reticularis***
  - Exit basal ganglion (GABA)
  - Evolutionary belongs to subtalamus

© David Kachlík 30.9.2015

Mid brain
Tegmentum
nuclei of cranial nerves

- ncl. **n. III** – at the level of colliculus superior
- ncl. **n. III accessorius dorsalis** *Edingeri-Westphali*
  - visceromotor (parasympatetic) nucleus for *mydriasis* and *accomodation*
- ncl. **n. IV** – at the level of colliculus inferior
- ncl. **mesencephalicus n. V**
  - Proprioceptive non migrated ganglion for eyeball and masticatory muscles

© David Kachlík 30.9.2015

Structure of n. III nucleus



© David Kachlík 30.9.2015

Mid brain
Tegmentum – other nuclei

- **substantia grisea centralis**
  = periakveductal grey (PAG)
  - By somatic and autonomic reactions deals with stress and pain
- **ncl. interstitialis *Cajali* + ncl. commissurae posterioris *Darkschewitzi***
  - From these originate fasciculus longitudinalis medialis by crossing at commissura posterior
  - Vertical eyeball movements
- **ncl. subbrachialis** = area ventralis tegmentalis *Tsai* (A10)
  - Part of RF
  - Produces dopamin for cortex and limbic system
- **ncl. interpeduncularis**
  - nucleus of limbic system

© David Kachlík 30.9.2015

Substantia grisea centralis (PAG)

Lateral PAG
defensive behavior
tachycardia
hypertension
non opioid analgesia



Ventrolateral PAG
quiet
hyporeactivity
bradycardia
hypotension
opioid analgesia

© David Kachlík 30.9.2015

Ncl. subbrachialis = area ventralis tegmentalis *Tsai* (A10)



© David Kachlík 30.9.2015

Mid brain
Tegmentum
tracts

- lemniscus
  - medialis, spinalis, trigeminalis, lateralis
  - Ascending tracts
- tractus tegmentalis centralis
- fasciculus longitudinalis medialis
- fasciculus longitudinalis posterior *Schützi*
- decussatio tegmentalis posterior (= dorsalis)
  - Crossing of tractus tectospinalis
- decussatio tegmentalis anterior (= ventralis)
  - Crossing of tractus rubrospinalis

© David Kachlík 30.9.2015

Mid brain
Tegmentum
Pedunculi cerebellares superiores

brachia conjunctiva

- After enter into tegmentum run in continuation of nucleus ruber
- On section at the level of colliculus inferior apparent as white caudal continuation
- Obsolete term „nucleus albus“
  - It is white matter !
- **decussatio pedunculorum cerebellarium superiorum**

  = full crossing close to nucleus ruber

  – tractus dentato-thalamo-corticalis

© David Kachlík 30.9.2015

Mesencephalon = Mid brain
Crus

only descending tracts

- **tractus pyramidalis (20 %)**
  - fibrae corticospinales (to alfa-motoneurons in spine)
  - fibrae corticonucleares (to nuclei of cranial nerves)
- **fibrae corticopontinae (80 %)**

  Run from cortex to nuclei pontis et arcuati, where get connected on tractus pontocerebellaris
  - fibrae frontopontinae *Arnoldi*
  - fibrae occipito,-temporo,-parietopontinae *Türcki*
- **fibrae corticoreticulares**
- **fibrae corticoolivares**

© David Kachlík 30.9.2015

Mesencephalic section at the level of COLLICULLUS SUPERIOR



1 - Colliculus superior
2 - Aquaeductus mesencephali */Sylvii/*
3 - Substantia grisea centralis
4 - Nucleus n. III.
5 - Tractus tectospinalis
6 - Decussatio tegmenti posterior
7 - Lemniscus medialis
8 - Nucleus ruber
9 - Tractus rubrospinalis
10 - N. III.
11 - Substantia nigra
12 - Fibrae frontopontinae */Arnoldi/* (Tractus coticopontinus)
13 - Fibrae corticonucleares (Tractus pyramidalis)
14 - Fibrae corticospinales (Tractus pyramidalis)
15 - Fibrae occipito-, parieto-, temporopontinae (Tractus corticopontinus)

Mesencephalic section at the level of COLLICULLUS INFERIOR



1 - Aquaeductus mesencephali + substantia grisea centralis
2 - Colliculus inferior
3 - Radix mesencephalica n. trigemini
4 - Lemniscus lateralis
5 - Lemniscus medialis
6 - Tractus tectospinalis
7 - "Nucleus albus"
8 - Deccussatio pedunculorom cerebellarium superiorum
9 - Substantia nigra
10 - Fibrae frontopontinae */Arnoldi/* (Tractus coticopontinus)
11 - Fibrae corticonucleares (Tractus pyramidalis)
12 - Fibrae corticospinales (Tractus pyramidalis)
13 - Fibrae occipito-, parieto-, temporopontinae (Tractus corticopontinus)
14 - Nucleus n. IV.
15 - N. IV.

© David Kachlík 30.9.2015

Unconsciousness and posture failure
Decortication

cut/lesion of stem rostrally to ncl. ruber

- desinhibition of tractus rubrospinalis
- desinhibition of tractus vestibulospinalis



© David Kachlík 30.9.2015

Unconsciousness and posture failure Decerebration

cut/lesion of stem caudally to ncl. ruber

example CMP of pons = locked-in syndrome



© David Kachlík 30.9.2015

Sectio mesencephali



A



B

A - tumor in aqueductu
B - hemiplegia alternans sup. (Weberi)
C - syndroma Benedikti



C

© David Kachlík 30.9.2015

syndroma Parinaudi

Hemiplegia alternans superior

Weber syndrom

<u>Tractus pyramidalis</u>

- contralateral hemiparesis
- central palsy of n. III

<u>N. III</u>

- ipsilateral ophthalmoplegia
- strabismus divergens
- mydriasis
- ptosis
- Acommodation loss
- Extinct light reflex



© David Kachlík 30.9.2015

Benedict syndrom

<u>lemnicus medialis</u>

- Contralateral hemianestesia

<u>nucleus ruber</u>

- Contralateral unvoluntary limb movements

Tumors close to aqueduct

- hydrocephalus internus

© David Kachlík 30.9.2015

Argyll-Ro

- pupilla: thin, no
  wide
- Convergence re
- *Reaction to ligh*
- Typical for syph
- Lesion of dorsa

Douglas Argyll-
1909) – scotish



© David Kachlík 30.9.2015

Parinaud syndrom

= dorsal mesencefalic syndrom

- **Palsy of looking up**
  - supranuclear lesion
  - Present reaction for „dolls head back eyes follow“, active look up is not possible x look down yes)
- **nystagmus**
  - konvergent-retractive: in attempt to look up ey retracts
- **Retraction of eyelid** (Collier sign)
- **Conjugated look down** („sign of setting s
- Often bilateral edema of discus (papilla) nervi optici
- *cause: pinealom, RS*

© David Kachlík 30.9.2015

Clinical syndromes in upper mid brain

A. Weber syndrom
B. Foix syndrom
C. Benedikt syndrom
D. Claude syndrom
E. Parinaud syndrom

1. tractus pyramidalis
2. pedunculus cerebellaris sup.
3. nucleus ruber
4. lemniscus medialis
5. colliculus superior
6. aqueductus mesencephali
7. fasciculus longitudinalis med.
8. nucleus n. III



© David Kachlík 30.9.2015

Injury

- Compression of n.III and n.IV
- incisura tentorii
- Herniation of part of temporal lobe when brain edema



© David Kachlík 30.9.2015

Further info

http://www1.indstate.edu/thcme/anderson/neurotext/neurotx1.html

© David Kachlík 30.9.2015